



# Pediatric Updates: Sepsis

Patrick Mahar, MD  
Section of Emergency Medicine  
Children's Hospital Colorado  
September 13, 2023



**Children's Hospital Colorado**

Affiliated with



University of Colorado  
Anschutz Medical Campus  
School of Medicine



Children's Hospital  
Colorado

# Financial Disclosures

- No relevant financial relationships with any commercial interests.

Patrick Mahar

# Objectives

- Review the physiologic aspects of sepsis and shock
- Discuss the differences between treating sepsis in adults and in kids (will keep this short!!!)
- Improve recognition/diagnosis of sepsis
- Review literature concerning sepsis in pediatrics
- Review the latest sepsis guidelines
- Discuss strategies for improving sepsis treatment/outcomes

# Sepsis

- Adult world:
  - Sepsis is bad.
  - Need to think about and recognize sepsis to treat sepsis
  - Recognizing sepsis early is very important
  - Avoid hypotension
  - Early antibiotics saves lives
- So, same in pediatric sepsis because they are all just little adults, correct?????



Children's Hospital  
Colorado

# Sepsis

- YES---they are basically little adults when it comes to sepsis!!!!
- Treatment of adult sepsis:
  - Rapid IVF, early antibiotics, avoid hypotension.
- Treatment of pediatric sepsis:
  - Rapid IVF, early antibiotics, avoid hypotension.





# Thank you

# Any Questions?



**Children's Hospital Colorado**

Affiliated with



University of Colorado  
Anschutz Medical Campus  
School of Medicine

Patrick Mahar, MD

[Patrick.mahar@childrenscolorado.org](mailto:Patrick.mahar@childrenscolorado.org)

# Pediatric Sepsis

- Pediatric Severe Sepsis
  - >75,000 yearly cases in US
  - US In-Hospital Mortality estimates
  - Over \$4.8 billion in healthcare costs in the United States



# Pediatric Definitions

- **Sepsis**
  - SIRS in the presence of infection
- **Severe Sepsis**
  - Sepsis + CV dysfunction OR ARDS OR  $\geq 2$  other organ dysfunction
- **Septic Shock**
  - Sepsis and CV organ dysfunction (hypotension, pressors use or elevated lactate)

## **Systemic Inflammatory Response Syndrome**

(2/4, 1 must be temp or wbc):

- Core Temp  $> 38.5^{\circ}\text{C}$  or  $< 36^{\circ}\text{C}$
- Tachycardia / Bradycardia
- Tachypnea
- WBC elevated or depressed



# Case #1

## 18 mo fever x 3 days

- 18 mo female; no PMHx
- Cough, congestion and runny nose for 5 days
- Fever started 3 days ago. T<sub>max</sub>=103.5
- Tylenol given but fever returns
- Seen by PCP 2 days ago and diagnosed with bronchitis and a double ear infection, started on Azithromycin
- Post-tussive emesis x4 over past 24 hours
- Decreased PO x1 day
  
- Immunizations: Has not received 18 mo shots yet
- Medications: Tylenol and azithromycin




Children's Hospital  
Colorado

# Case #1

## 18 mo fever x 3 days

- T: 40.1 HR: 191 RR: 40 Pox: 92% on RA
- Clinging to MOC; sleepy but cries when you approach
- Nose: Copious amounts of clear/yellow nasal drainage
- Ears: TM: erythematous bilat. Clear fluid
- Lungs: course BS/crackles bilat
- Heart: S1S2 Tachy



# Case #1

## 18 mo fever x 3 days

- T: 40.1 HR: 191 RR: 40 Pox: 92% on RA **BP: 82/48**
- ARE WE IN SHOCK????
- What do you want to do?
  - IVF, LABS, Abx
  - OR
  - Motrin/Tylenol and re-eval



Children's Hospital  
Colorado

# What is shock?

- What is shock?
- What vital signs define shock?
- What physical exam findings are consistent with shock?



Children's Hospital  
Colorado

# SHOCK

**OXYGEN SUPPLY FAILS TO MEET OXYGEN DEMAND**

OXYGEN  
SUPPLY



OXYGEN  
DEMAND



## OXYGEN DELIVERY

CARDIAC OUTPUT X ARTERIAL OXYGEN CONTENT

Cardiac  
Output

Heart rate

-Intravascular  
volume  
-SVR

Stroke Volume

Preload  
After load  
Contractility

Arterial oxygen content

-Hemoglobin  
-Oxygen Saturation  
-Partial pressure of  
oxygen dissolved in  
plasma

# SHOCK

- Compensated Shock

- **Normal blood pressure**
  - Body is maintaining blood pressure in normal range
    - ↑ HR and SVR to maintain BP
- Early shock
- Need to recognize and act to prevent progression


- Uncompensated Shock

- **Hypotension**
- Late finding
- BAD SIGN → BIG TROUBLE
- “Assume the crash position”



# Vital signs: When to worry

- $BP = \text{Cardiac Output} \times \text{System Vascular Resistance (SVR)}$
- $CO = \text{Stroke Volume} \times \text{Heart Rate}$
- Quick Estimate of Hypotension in Pediatric Patient
- $SBP \leq (2 \times \text{Age}) + 70$ 
  - 4 yo
  - $SBP \leq (2 \times 4) + 70$
  - $SBP \leq 78$







## Case #2 R.S.

- 12y.o male No PMhx.
- Presents to PCP with 1 day of fever, leg pain and vomiting.
- Reports throwing up the first time during the night and then again in AM. MOC reports was crying his legs hurt and he had trouble walking from car to office b/c of legs hurting.
- Tmax=104 this AM
- 2 days ago dove for loose ball while playing basketball but didn't hurt legs, only small cut on right arm.
- Pt throws up in office waiting room and again when throat swap was obtained.
- T=102 HR=140 RR=36



# Case #2 R.S.

- Exam notable for:
  - Gen: Ill/weak appearing
  - Mucous membranes-dry
  - Neck: supple FROM
  - Abd: soft, NT, ND, no masses.
  - Ext:
    - Right upper extremity with 1 cm laceration w/ scab in place minimal erythema and no red streaks up arm
    - Lower Ext: bilateral tenderness of thighs R>>L; FROM at knees
  - Skin- mottled appearance and delayed cap refill
- Rapid strep negative
- NOW WHAT???



## Case #2 R.S.

- PCP tells family to go to the academic hospital nearby for IVF to make him feel better. Tells family he has stomach flu and will likely have diarrhea to go along with this.
- 19:14 arrived at hospital ED:
- HR: 143 RR: 28 T:100.9 BP 118/48 Wt: 77kg
- Pt treated with Zofran, 2 liters NS bolus and labs sent.
- 21:14 Pt no longer vomiting, thus decision made to d/c
- 21:26 T: 102 HR: 131 RR: 22 BP: 108/42
- D/c home with dx of stomach flu;
- Plan: Supportive care at home
- 00:30: Labs printed off showing ↑ WBC, (marked left shift)



## Case #2 R.S.

- Pt continues to have fever and emesis during the night.
- Call PCP in AM instructed to alternate Tylenol and Motrin for fever.
- Pt develops diarrhea.
- Call PCP again and MOC describes patient as weak. Instructed to have him drink and eat some crackers.
- Pt screams in pain when legs are touch and is too weak to sit up
- MOC calls PCP and says his nose and legs are turning blue
- PCP instructs family to return to ED.



Children's Hospital  
Colorado

# Case #2 RS

- Admitted to ICU
  - Intubated
  - Renal failure
  - DIC
  - Codes x2 and brought back
  - Day 3 arrest 3<sup>rd</sup> time and dies.
- 
- Group A strep sepsis



Children's Hospital  
Colorado

- 2012 Death of Rory Staunton from unrecognized Group A Strep sepsis
- "Rory's Law"
- New York State legislates pediatric sepsis protocol requirements



The New York Times

N.Y. / Region

WORLD | U.S. | N.Y. / REGION | BUSINESS | TECHNOLOGY | SCIENCE | HEALTH | SPORTS

T-W-I-D-D-Y  
K-N-O-W-S  
O-C-E-A-N-F-R-O-N-T...297 HOMES



## Cuomo Plans New Rules in Fight Against Sepsis

By JIM DWYER

Published: January 7, 2013

# SEVERE SEPSIS TRIAGE SCREENING TOOL

RORY'S RESULTS

Does the patient have any three of the following?

- Suspected Infection
- Temp > 100.4 or < 96.5 or rigors
- HR > 90
- RR > 20
- Any alteration of mental status
- O2 Sat < 90%
- SBP < 90
- Suspected/Known Immunocompromise (AIDS/Active Cancer/Organ Transplant Patient)

Yes  No

*If Yes clicked:*

Go to Nursing Sepsis Panel Orders.

### Triage Sepsis Panel Orders

(all boxes should be checked by default)

- Notify clinician to initiate verbal order for sepsis panel
- CBC
- Metabolic Panel
- Lactate (venous or arterial)
- Draw and Hold PT/PTT
- Draw and Hold Blood Cultures
- Record Vital Signs Q1 hour plus temperature

Is the patient's SBP < 90 or MAP < 65

Yes  No

*If Yes clicked:*

Present case to physician.

*If patient meets criteria, but you feel the patient doesn't need lab testing, please speak to*



NYU Hospitals Center  
Patient Record  
NYUMC

Patient: STAUNTON,  
DOB: 05/13/1999  
Attending: [REDACTED]  
Admit Date: 03/29/2012  
Location: EMERG-TH

This report contains documentation entered between 03/29/2012

### Vital Signs & Measurements F/S General

03/29/2012 21:26

Authored By:

#### BLOOD PRESSURE

Blood Pressure Systolic Systolic : 103 mmHg  
Blood Pressure Diastolic Diastolic : 50 mmHg  
Blood Pressure Mean Mean : 67 mmHg

#### TEMPERATURE

Temperature Temperature (F) degree F : 102 degrees F  
Temperature Temperature (C) degree C : 38.8 degrees C

#### PULSE

Pulse Pulse Rate Rate (bpm) : 131 bpm

#### RESPIRATORY

Respirations Respiratory Rate Rate /min : 22 /min

#### PULSE OXIMETRY

Pulse Oximetry Saturation O2 Sat (room air) % (Room Air) : 99  
29-Mar-2012 21:26 Vital Signs & Measurements F/S General  
Norma (Nursing Attendant, Nursing Attendant, BENNEN02)

# Sepsis Kills.

**250,000 Americans die each year from sepsis.** That's more than from **AIDS, breast cancer and prostate cancer COMBINED.** Sepsis is the body's life-threatening reaction to an infection. Anyone can get sepsis. A small cut, a bug bite or an infected tooth can all lead to sepsis.

Sepsis is preventable and treatable.  
**Do you know the Signs of Sepsis?**



Fever/  
Shivering or  
Very Cold



Rapid  
Breathing



Extreme Pain/  
Physical  
Discomfort



Pale or  
Mottled  
Skin



Disoriented/  
Confused &  
Sleepy/Difficult  
to Wake



Elevated  
Heart Rate

THE RORY STAUNTON FOUNDATION

FOR SEPSIS PREVENTION

**Help Save Lives. Share the Signs of Sepsis with your family and friends.**

For more information, visit [www.rorystauntonfoundationforsepsis.org](http://www.rorystauntonfoundationforsepsis.org)





Children's Hospital  
Colorado

# How do we do better???

- Early Recognize of sepsis (and those at risk)
  - Rapid IV Access
  - Aggressive Fluid Resuscitation
  - Early Antibiotics
  - Reassess – **DO NOT TOLERATE HYPOTENSION**
  - Early use of pressors
- 
- Sepsis pathways and order bundles save lives!!!



Children's Hospital  
Colorado

# www.childrenscolorado.org



[Careers](#)

[Healthcare Professionals](#)

[Contact](#)

[Locations](#)

[MyChart](#)

[Schedule an Appointment](#)

[Donate](#)



Children's Hospital Colorado

[Doctors & Departments](#)

[Conditions & Advice](#)

[Your Visit](#)

[Community](#)

[Research & Innovation](#)



#1 in Colorado

#1 in the region

Our personalized care leads to miraculous things. That's why we're ranked among the best children's hospitals in the nation by U.S. News & World Report.

*Here, it's different.™*



# Clinical Pathways



delivering the best possible treatment across the continuum of care.

[Directory of Services](#)

[PedsConnect](#)

[Physician Relations](#)

[Lab and Microbiology Test Directory](#)

[Pediatric nursing resources](#)



Referral tools



Clinical pathways



Continuing medical education

## Healthcare Professionals

[COVID-19 Resources](#)

[Clinical Resources](#)

**2000+**

Board-certified pediatric  
specialists

**TOP 5**


Research hospital in the U.S.

**TOP 25**

In the U.S. for all 10 specialties

# Sepsis Pathway: Initial Management

## ALGORITHM 1. ED/UC SEPSIS



**Quick Links**  
[Emergency Abx Selection for Sepsis STAT](#)  
[Urgent Access Algorithm](#)





Children's Hospital  
Colorado

# Early Recognition

- How do we know when to be concerned for sepsis
- Can't treat sepsis without thinking of sepsis
-



# What Clinical Signs and Hemodynamic Variables Can be Used to Direct Treatment of Newborn and Pediatric Shock?

Table 2. American College of Critical Care Medicine hemodynamic definitions of shock

Cold or warm shock

Decreased perfusion manifested by altered decreased mental status, capillary refill  $>2$  secs (cold shock) or flash capillary refill (warm shock), diminished (cold shock) or bounding (warm shock) peripheral pulses, mottled cool extremities (cold shock), or decreased urine output  $<1$  mL/kg/h



Children's Hospital  
Colorado





Children's Hospital  
Colorado





# Physical Exam for Detection

**Table 2 Sepsis clinical recognition signs present in ED as predictors of organ dysfunction within 24 hours**


Individual predictor	Prevalence n (%)	Sensitivity	Specificity	Negative predictive value	Positive predictive value	Positive likelihood ratio	Concordance**
Altered Mental Status	43 (18%)	0.54 (0.29–0.77)	0.84 (0.78–0.88)	0.97 (0.93–0.99)	0.16 (0.07–0.30)	3.3* (1.8–5.9)	76%
Abnormal Capillary Refill	36 (15%)	0.08 (0.01–0.33)	0.85 (0.79–0.89)	0.94 (0.90–0.97)	0.03 (0.001–0.15)	0.5 (0.1–3.4)	96%
Abnormal Peripheral Pulses	8 (3%)	0.15 (0.04–0.42)	0.97 (0.94–0.99)	0.95 (0.92–0.98)	0.25 (0.03–0.65)	5.8* (1.2–26.0)	92%
Cold/Mottled Extremities	5 (2%)	0.08 (0.01–0.33)	0.98 (0.95–0.99)	0.95 (0.91–0.97)	0.20 (0.01–0.72)	4.3 (0.5–36.2)	100%
<b>Number of predictors</b>							
≥ 1	77 (32.2%)	0.62 (0.32–0.86)	0.69 (0.63–0.75)	0.97 (0.93–0.99)	0.10 (0.05–0.19)	2.0* (1.3–3.2)	72%
≥ 2	15 (6.3%)	0.23 (0.05–0.54)	0.95 (0.90–0.97)	0.96 (0.92–0.98)	0.20 (0.04–0.48)	4.4* (1.4–13.5)	72%

\*Statistically significant associations.

\*\*Based on 25 patients with two independent assessments.

# Sepsis Pathway: Initial Management

## ALGORITHM 1. ED/UC SEPSIS



# Sepsis = Infection + Organ Dysfunction

- Many organ dysfunctions require laboratory testing
- Diagnosis does not require microbiological confirmation
- Start treatment for suspicion of infection + organ dysfunction
- Consider de-escalation if clinical picture/labs do not support infection

CV: Hypotension, lactate, vasopressor use

Resp: New positive-pressure ventilation requirement


Neuro: Altered mental status

Renal: Acute kidney injury ←

Hepatic: Elevated LFTs ←

Heme: DIC, low platelets ←

Require labs to determine, thus takes time, so act quickly if have other signs.



- Verbal or paging notification per site protocol
- In NOC, begin request for transport to Anschutz
- Use ED Sepsis STAT order set

**1. Activate & Notify Say "Sepsis"**

- Verbal or paging notification per site protocol
- In NOC, request transport to Anschutz as soon as a need for higher level of care develops
- Use ED Sepsis Yellow order set

**IV Access x2**

- 2 peripheral IVs with largest gauge possible and/or central access
- Urgent Access Algorithm for rapid access

**2. Rapid Access & Labs**

**PIV or Central Line Access Immediately**

- Urgent Access Algorithm for rapid access

**Labs per Order Set**

- Cultures, CBC, lactate, VBG, CMP, DIC panel

**Labs per Order Set**

- Cultures, CBC, lactate

**Bolus**

- 10-20 mL/kg isotonic crystalloid (push-pull, pressure bag or rapid infuser for STAT)
- Upgrade to STAT if bolus rate faster than 1 hour or vasopressor required

**Antibiotics**

- STAT: complete antibiotics coverage immediately per order set
- Yellow: expedited antibiotics if clinically indicated

**3. Resuscitation**  
Consider bundle steps simultaneously

**Vasopressors**

- Within 60 minutes of first hypotension
- Do NOT tolerate hypotension, consider epinephrine or norepinephrine

**Reassess**


- Repeat until clinical euvoemia
- Vital signs, serial lactate, POCUS, CVP where available

**! Escalate to STAT for:**

- Decompensation
- Hypotension
- Organ dysfunction




# Why is organ dysfunction important?



# Lactate and Sepsis

- Lactate in recognition and treatment in adult sepsis is well studied and accepted practice
  - Lactate is associated with increased mortality in adults w/sepsis
  - Lactate clearance correlates with improved outcomes
  - Use of lactate and lactate clearance to direct therapy is associated with equal or superior outcomes for adults with sepsis
  - Clinical care bundles that include use of lactate have shown to improve outcomes.
- What does elevated lactate mean for the patient?

# 5x Greater Risk



- Verbal or paging notification per site protocol
- In NOC, begin request for transport to Anschutz
- Use ED Sepsis STAT order set

**1. Activate & Notify  
Say "Sepsis"**

- Verbal or paging notification per site protocol
- In NOC, request transport to Anschutz as soon as a need for higher level of care develops
- Use ED Sepsis Yellow order set

- IV Access x2**
- 2 peripheral IVs with largest gauge possible and/or central access
  - Urgent Access Algorithm for rapid access

**2. Rapid Access  
& Labs**

- PIV or Central Line Access Immediately**
- Urgent Access Algorithm for rapid access

- Labs per Order Set**
- Cultures, CBC, lactate, VBG, CMP, DIC panel

- Labs per Order Set**
- Cultures, CBC, lactate

- Bolus**
- 10-20 mL/kg isotonic crystalloid (push-pull, pressure bag or rapid infuser for STAT)
  - Upgrade to STAT if bolus rate faster than 1 hour or vasopressor required

- Antibiotics**
- STAT: complete antibiotics coverage immediately per order set
  - Yellow: expedited antibiotics if clinically indicated



**3. Resuscitation**  
Consider bundle steps simultaneously

- Vasopressors**
- Within 60 minutes of first hypotension
  - Do NOT tolerate hypotension, consider epinephrine or norepinephrine

- Reassess**
- Repeat until clinical euvoemia
  - Vital signs, serial lactate, POCUS, CVP where available

**! Escalate to STAT for:**

- Decompensation
- Hypotension
- Organ dysfunction







Children's Hospital  
Colorado

# Fluids





Chi



Pull from IV bag

Turn stopcock

Push to patient

← Repeat →



Children's Hospital  
Colorado

# Rapid Infuser

- Flow rates from 2.5-1000ml/min
- Temperature controlled fluids
- Can be used down to 10kg.





Children's Hospital  
Colorado

# LifeFlow



- Rapid infusion of fluids and/or blood
- Deliver 500ml of fluid in less than 2 minutes through a 20G IV catheter
- 4x faster than pressure bag for fluids and ~3x faster for blood

# First evidence for early fluid in pediatric sepsis

- All children with septic shock with PA catheter by 6 hours
- 34 patients, mean age 13.5 months






	<b>1<sup>st</sup> Hour Fluid</b>	<b>N</b>	<b>Mortality</b>
Group 1	<20 ml/kg	14	57%
Group 2	20-40 ml/kg	11	64%
Group 3	>40 ml/kg	9	11%

- ARDS (n=11), cardiogenic pulmonary edema (n=5) not associated with volume received
- At time of PA placement: Hypovolemia more frequent in Groups 1&2, all hypovolemic patients died (n=8)




# How much fluid should I give?

- In most cases, start with at least 20 ml/kg, some patients may require up to 40-60 ml/kg or more to achieve euvolemia.
- If you have achieved euvolemia, per your clinical assessment, and the patient remains hypotensive, start vasopressors/inotropes.
- There is debate about exactly what amount of fluid is right for sepsis, and it is probably different for each child.
  - If child doesn't have a specific fluid sensitivity (such as a brain tumor, renal failure, heart failure or severe anemia)
  - Being studied but no definitive studies yet.



ALGORITHM 3. EMERGENT ANTIBIOTIC SELECTION FOR SEPTIC SHOCK/SEPSIS STAT





# ALGORITHM 3. EMERGENT ANTIBIOTIC SELECTION FOR SEPTIC SHOCK/SEPSIS STAT


Suspected Infection and Critical Illness  
Acute Organ Dysfunction / Shock / Sepsis STAT

**Inclusion Criteria:**  
Initial antibiotics for:

- Suspected infection and critical illness
- ED Sepsis STAT patients
- Inpatients with sepsis plus acute organ dysfunction and/or shock

**Exclusion Criteria:**

- Age less than 60 days



Anaerobe

Suspect anaerobe? (e.g. sinus, abdominal source, Lemierre's)

Yes

Metronidazole

No

Toxin

Suspect toxin-mediated infection? (e.g. toxic shock, skin/soft tissue source)

Yes


Clindamycin

No

Initial antibiotic algorithm complete; consult ID/subspecialist for complex infections/history

### Additional Considerations:

- For suspected meningitis, consider dexamethasone before or shortly after first antibiotic dose.
- Consider ID consultation for allergies, history of resistance (MRSA), yeast coverage or other complicating factors. Primary service should be consulted as soon as feasible.
- For CNS coverage, ceftriaxone may require dosing every 12 hours. Please discuss at handoff.
- Check prior +cultures.
- Ongoing care team should review and continue antibiotics as indicated.



- Verbal or paging notification per site protocol
- In NOC, begin request for transport to Anschutz
- Use ED Sepsis STAT order set

**1. Activate & Notify Say "Sepsis"**

- Verbal or paging notification per site protocol
- In NOC, request transport to Anschutz as soon as a need for higher level of care develops
- Use ED Sepsis Yellow order set

**IV Access x2**

- 2 peripheral IVs with largest gauge possible and/or central access
- Urgent Access Algorithm for rapid access

**2. Rapid Access & Labs**

**PIV or Central Line Access Immediately**

- Urgent Access Algorithm for rapid access

**Labs per Order Set**

- Cultures, CBC, lactate, VBG, CMP, DIC panel

**Labs per Order Set**

- Cultures, CBC, lactate

**Bolus**

- 10-20 mL/kg isotonic crystalloid (push-pull, pressure bag or rapid infuser for STAT)
- Upgrade to STAT if bolus rate faster than 1 hour or vasopressor required

**Antibiotics**

- STAT: complete antibiotics coverage immediately per order set
- Yellow: expedited antibiotics if clinically indicated

**3. Resuscitation**  
Consider bundle steps simultaneously

**Vasopressors**


- Within 60 minutes of first hypotension
- Do NOT tolerate hypotension, consider epinephrine or norepinephrine

**Reassess**

- Repeat until clinical euvolemia
- Vital signs, serial lactate, POCUS, CVP where available

**! Escalate to STAT for:**

- Decompensation
- Hypotension
- Organ dysfunction




# Mortality and functional morbidity after use of PALS/APLS by community physicians

- Early shock reversal was associated with reduced mortality (5.06% vs 16.37%) and functional morbidity (1.56% vs 4.11%) rates.
- Early use of PALS/APLS-recommended interventions was associated with reduced mortality (8.69% vs 15.01%) and functional morbidity (1.24% vs 4.23%) rates.

## **CONCLUSIONS:**

- Pediatric shock recognition and resuscitation in the community hospital improves survival and functional outcome regardless of diagnostic category.
- The development of shock/trauma systems for children with and without trauma seems prudent.

# Mortality in patients referred for shock for transport to pediatric ICU



**FIGURE 1**

Mortality rates increase according to the degree of hemodynamic abnormality at presentation to the community hospital. CR indicates capillary refill time. <sup>a</sup>  $P < .05$  versus no shock.



Children's Hospital  
Colorado

# Early Reversal of Shock

- At arrival of transport team (median 75 min): 26% achieved shock reversal – associated with 96% survival
- Each additional hour persistent shock...

Han YY, Carcillo JA, Dragotta MA, Bills DM et al. Early reversal of pediatric-neonatal septic shock by community physicians is associated with improved outcome. *Pediatrics*. 2003;112:793-799.



Children's Hospital  
Colorado

# Early Reversal of Shock

- At arrival of transport team (median 75 min): 26% achieved shock reversal – associated with 96% survival
- Each additional hour persistent shock...

## Risk of death doubled

Han YY, Carcillo JA, Dragotta MA, Bills DM et al. Early reversal of pediatric-neonatal septic shock by community physicians is associated with improved outcome. *Pediatrics*. 2003;112:793-799.



# Treat hypotension

- IVF bolus-
  - 20 ml/kg
  - Give rapidly
  - Reassess/Repeat
  - Goal improved circulation (~60 ml/kg) in first 60 min
- Pressors:
  - Epi/NorEpi (yes can be given through PIV)
- Steroids:
  - If concern for adrenal insufficiency → Hydrocortisone
    - Infants: 25mg
    - Children: 50mg
    - Adolescents: 100mg





# Do Not Tolerate Hypotension

- Traditionally learned to give all fluid first (~60ml/kg), and then start pressors, it may be better to work on both simultaneously (try to get at least 1-2 boluses in first).  
*(Shrink the tank and fill it at the same time).*
- The new pediatric sepsis guidelines recommend that no patient should be hypotensive longer than 1 hour without having a vasopressor/inotrope on board.
- If the patient has been hypotensive for 1 hour, we want to see pressors on board, so please start ordering and planning for them by the 45 minute mark.



Children's Hospital  
Colorado

# First-Line Vasoactive Agents (an ED guide)

	Inotropy	Systemic Vascular Resistance	BP	Use in Sepsis?
Norepinephrine (alpha > beta)	+	+++	↑	Most patients Premixed at all CHCO sites, More familiar in general ED's
Epinephrine (beta > alpha)	+++	+	↑	Younger Community-acquired gram positive Significant cardiogenic component
Dopamine (5-10 mcg/kg/min) Beta > alpha, dopa	++	++	↑	If desperate Problems: 'dirty drug' (variable effects at different doses), arrhythmogenic, HPA/immune effects (decreases GH, prolactin)

Vasopressin, milrinone, dobutamine... potentially useful, most sepsis should be in ICU by then...

# Epi/NorEpi

- Can be given peripherally in an emergency, but central access is preferred.
  - There have been significant instances of patient harm from infiltration (KEEP CLOSE EYE ON IV)
  - This is even more of a concern in the early phases of resuscitation when our IV's (and IO) may be tenuous and patients may be moved and transported.
- Be careful with dosing.
- Let pharmacy make drips whenever possible.



# Sepsis Treatment And Recognition Program

## Children's Hospital Colorado



Children's Hospital Colorado

Affiliated with



University of Colorado  
Anschutz Medical Campus  
School of Medicine

# Sepsis Stat

**Does the patient have fever and/or concern for infection with any of the below symptoms:**

- **Hypotension with tachycardia or poor perfusion?**
- **Severe alterations in mental status/perfusion/respiratory distress?**
- **Critically-ill appearance?**

# Sepsis Yellow

**Does the patient have fever with concern for infection with any of the below symptoms:**


- Persistent/worsening tachycardia despite treatment of fever & dehydration?
- Immunosuppression/immuno-deficiency or central line and worsening clinical status?
- Consider for:
  - Changes to mental status + infection
  - Cool extremities, poor capillary refill, diminished pulses, mottling
  - Flushed, warm extremities, bounding pulses, flash capillary refill



Children's Hospital  
Colorado

# Sepsis Activation

- Verbal request to unit secretary
  - Nursing or physician can initiate a sepsis activation
- Secretary activates page to relevant staff:
  - Charge nurse (ED / ICU), Pharmacist, Physicians
- Secretary clicks a timestamp button in EPIC




# Order Sets

## Pre-checked:


Vitals  
Fluid  
Laboratories

## Key antibiotic elements:

- Indication in order set
- Dosing preprogrammed
- Max dose automatic correction
- Can be ordered in one "click"




# ED Sepsis Stat Patients: Percent Receiving Antibiotic in 60 min of Triage





# Outcomes: Hospital Length of Stay





Children's Hospital  
Colorado

## Case #3

- 18:52 Phone call: 4 year old won't drink and has a fever. Language barrier.
- 19:10 EMS arrival on scene
- 4 year-old, no PMH
- Seen 2 weeks ago at ED for fever and was given medicine that made him better
- Recovered, was back at pre-school
- Now 4 days of new fever, worsening cough, vomiting and not drinking
- Mom reports: Has only had 1 can Mountain Dew for 24 hours.. No urine output in 12 hours



Children's Hospital  
Colorado

# Case#3: On Scene

- 19:12 T=100 HR:132 RR:30 SpO2: 86%
- Moaning and grabbing abdomen
- Crying, moaning, grunting
- Refuses to walk – carried to stretcher
  
- 19:25 Call to Hospital:
  - “This is \*\*\*\* agency coming to ED non-emergent with a 4 year old male for fever and vomiting. Vitals are HR: 148 RR:30 Pox: 88% but I don’t believe it bc hands are cold and not getting good reading, but he is on blow-by O2. We are 10 minutes out.”



Children's Hospital  
Colorado

# Case #3 in ED

- 19:39 Arrive at ED → Room 12
  - VS: T-102 RR:34 BP: 82/42 Pox: 86% on RA Wt: 15 kg
- 19:52 Orders:
  - Chest xray
  - Zofran
  - Motrin
- 21:30 CXR shows possible LLL pneumonia
  - PO amoxicillin ordered
- 21:45
  - VS T-100.6 HR:164 RR: 60 BP: 74/32 Pox=92 on 4-5L NC,
  - Sleepy; does not cry with exam
  - Extremities cool, weak pulses
- 21:48 Provider alerted to change in status



Children's Hospital  
Colorado

## Case #3 in ED

- 21:50 IVF bolus, CBC, Blood cultures ordered
- 21:52 IV attempt R arm not successful
- 21:53 IV attempt R hand not successful
- 21:55 IV attempt L hand not successful
- 21:58 IV L foot not successful
- 22:04 Charge nurse to bedside for IV attempt
- 22:06 22g in right foot.
  - Labs obtained/sent.
  - IVF 20ml/kg started at 300ml/hr
- 22:32 Pt unresponsive.
- 22:35 VS: HR 164 BP:63/23 RR: 12. Intubation
- 22:42 CPR started



Children's Hospital  
Colorado

# Case #3 in ICU

- Day 1-
  - NorEpi drip
  - No urine output
  - DIC
- Day 2-
  - CPR required x2
  - No brain activity on EEG
- Day 3
  - Family removes support and pt passes.
- Autopsy: Group A Strep grew from pulmonary fluid  
+Influenza



Children's Hospital  
Colorado

# What went wrong?

- What were the warning signs?
- What were the reassuring signs?
- What steps could have been better?



Children's Hospital  
Colorado

# What do you notice about this case?

- Initial vitals not that bad
- Exam findings may be subtle
- Return of fever, or worsening, after initial febrile prodrome
- No focus on blood pressure, early access, IV fluid, antibiotics in the treatment plan until too late
- High-risk difficulty airway for intubation





Children's Hospital  
Colorado

# What makes an intubation a difficult intubation?

- Trauma- Direct airway trauma and/or C-collar
- Anatomical abnormalities
- Inexperience for patient type
- Hemodynamically unstable patient
  - Hypotension
  - Hypoxia
  - Acidosis

# Risk Factors for Peri-intubation Cardiac Arrest in a Pediatric Emergency Department

*Nicholas Pokrajac, MD,\* Emily Sbiroli, MD,† Kathryn A. Hollenbach, PhD, MPH,‡ Michael A. Kohn, MD, MPP,\* Edwin Contreras, MD,§ and Matthew Murray, MD†* PEC Jan 2022

**TABLE 2.** Hemodynamic, Respiratory, and Intubation Characteristics of Cases and Controls

	PICA (n = 21)	Controls (n = 84)	OR (95% CI)	P
Hemodynamic and respiratory characteristics				
Elevated HR	11 (52.4)	53 (63.1)	0.6 (0.2–1.7)	0.455
→ Systolic hypotension (or unobtainable)	12 (57.1)	6 (7.1)	<b>17.3 (5.2–57.5)</b>	<b>&lt;0.001</b>
→ Diastolic hypotension (or unobtainable)	11 (52.4)	6 (7.1)	<b>14.3 (4.3–47.1)</b>	<b>&lt;0.001</b>
Elevated SI	6 (37.5)	17 (20.2)	2.4 (0.8–7.4)	0.191
→ Delayed CRT (>2 s)	18 (85.7)	19 (22.6)	<b>20.5 (5.5–77.2)</b>	<b>&lt;0.001</b>
Received at least 10 mL/kg IVF	5 (23.8)	31 (36.9)	0.5 (0.2–1.6)	0.312
→ Hypoxia (or unobtainable)	13 (61.9)	2 (2.4)	<b>66.6 (12.7–349.1)</b>	<b>&lt;0.001</b>



**TABLE 2.** Hemodynamic, Respiratory, and Intubation Characteristics of Cases and Controls

	PICA (n = 21)	Controls (n = 84)	OR (95% CI)	P
Type of paralytic agent				0.639
Rocuronium	13 (61.9)	72 (85.7)		
Vecuronium	2 (9.5)	7 (8.3)		
Succinylcholine	1 (4.8)	2 (2.4)		
→ No paralytic agent	5 (23.8)	2 (2.4)	<b>12.8 (2.3–71.9)</b>	<b>0.003</b>
Type of sedative agent				0.452
Etomidate	3 (14.3)	29 (34.5)		
Ketamine	2 (9.5)	5 (6.0)		
Benzodiazepine	2 (9.5)	24 (28.6)		
Pentobarbital	3 (14.3)	16 (19.0)		
Other	1 (4.8)	7 (8.3)		
→ No sedative agent	10 (47.6)	3 (3.6)	<b>24.5 (5.8–103.2)</b>	<b>&lt;0.001</b>
Night intubation (7:00 PM to 7:00 AM)	9 (42.9)	29 (34.5)	1.4 (0.5–3.8)	0.612
→ Greater than 1 intubation attempt	14 (66.7)	23 (27.4)	<b>5.3 (1.9–14.8)</b>	<b>0.001</b>




Children's Hospital  
Colorado

ORIGINAL CONTRIBUTION



# Identification of the Physiologically Difficult Airway in the Pediatric Emergency Department

Preston N. Dean, MD<sup>1,2</sup> , Erin F. Hoehn, MD<sup>1,2,3</sup>, Gary L. Geis, MD<sup>1,2</sup>, Mary E. Frey, MSN<sup>1</sup>, Mary K. Cabrera-Thurman<sup>1</sup>, Benjamin T. Kerrey, MD, MS<sup>1,2</sup>, Yin Zhang<sup>4</sup>, Erika L. Stalets, MD, MS<sup>2,5</sup>, Matthew W. Zackoff, MD, MEd<sup>2,5</sup>, Andrea R. Maxwell, MD<sup>2,5</sup>, Tena M. Pham<sup>5</sup>, and Andrew J. Lautz, MD<sup>2,5</sup>

ACADEMIC  
EMERGENCY  
MEDICINE 2020

-primary outcome: peri-intubation cardiac arrest, defined as cardiac arrest (documented chest compressions or non-perfusing rhythm within 10 minutes of the completion of tracheal intubation)



Ch Table 1  
Clinical Characteristics of Patients Undergoing Tracheal Intubation in the PED by Risk Group

Characteristic	High Risk ( <i>n</i> = 36)	Standard Risk ( <i>n</i> = 177)	p-value*
Age (months)	8.5 (1-118.5)	18 (2-89)	0.489
Indication for RSI			<0.0001
Respiratory	16 (44)	71 (40)	
Altered mental status	9 (25)	30 (17)	
Sepsis	5 (14)	0	
Cardiac arrest	3 (8)	0	
Seizure	3 (8)	76 (43)	
High-risk criteria met			Not applicable
Hypotension	15 (42)	0	
Cardiac dysfunction	12 (33)	0	
Hypoxia	11 (31)	0	
Metabolic acidosis	8 (22)	0	
Post-ROSC	7 (19)	0	
Status asthmaticus	2 (6)	0	

Table 2  
Outcome Data for Patients Undergoing Tracheal Intubation in the PED by Risk Group

Outcome	High Risk ( <i>n</i> = 36)	Standard Risk ( <i>n</i> = 177)	% Difference (95% CI)	p-value*
Peri-intubation arrest	2 (5.6)	0	5.6 (1.0 to 18.1)	0.0279
Any postintubation arrest in STS	4 (11.1)	0	11.1 (4.1 to 25.3)	0.0007
In-hospital mortality	9 (25)	4 (2.3)	22.7 (11.0 to 38.9)	<0.0001
ECMO	3 (8.3)	0	8.3 (2.5 to 21.8)	0.0044
First-attempt success	17 (47.2)	117 (66.1)	-18.9 (-35.5 to -1.5)	0.0382

Data are reported as *n* (%).



Children's Hospital  
Colorado

# Physiologically Difficult Airway

Patients at increased risk of peri-intubation decompensation if have:

- Hypoxia
- Hemodynamic instability
- Cardiac dysfunction
- Acid/base derangements
- Asthma patient



Children's Hospital  
Colorado

# Plan for Successful RSI

1. Identifying the physiologically difficult
2. Do you need to intubate? Do we need to **RIGHT NOW?**
3. Optimization of physiologic status prior to tracheal intubation
  1. Correct Hypoxia— pre-oxygenation, apneic oxygenation (don't tolerate hypoxia)
  2. Get as close as possible to hemodynamically stable
    1. IVF bolus (improve preload)
    2. Pressors (improve cardiac squeeze and vascular tone → preload)
      1. Epi drip, or “Low dose Epi” (1/10<sup>th</sup> code dose epi)
  3. Correct acidosis (Myocardium doesn't like acidosis)
4. Having cardiac arrest precaution measures in place
5. Optimize first pass success- paralytic, induction agent, and most-experienced intubator



Children's Hospital  
Colorado

# Case#4

- 00:15 911call: 5 year-old, fever, not drinking
- 00:27 Arrive on scene:
  - 5 year-old, no PMH
  - Fever, malaise, cough x 1 day
  - Decreased po, no UOP today
- T=37.9 HR=146 RR=40 Sat 82% BP = 86/55
  - Sleepy but cries with exam (no tears) Coughing;
  - Heart:S1S2 Tachy
  - Lungs: marked decreased BS on left
  - Cap refill~4sec
- 00:42 O2 15 liters via NRB





Children's Hospital  
Colorado

# Case#4: Back of ambulance

- 00:45 Repeat Vitals
  - HR: 172 RR: 42 BP: 82/32 Pox:92% O2 15 liters via NRB
  - Broselow: 18kg
- 00:47 Call to ED:
  - "Agency \*\*\* coming to ED with 5 y.o with concern for septic shock. Pt with fever and poor PO intake and cough for 1 day. Current vital signs: HR 178 RR42 BP:80/32 Pox: 93%on 15 liters NRB. Pt is 18kg white by Broselow. Again I am concerned for septic shock and we are 8 minutes out"



Children's Hospital  
Colorado

# Case#4 in ED

- 00:48:
  - Trauma/resuscitation Room 1 prepared
    - Nursing staff, respiratory therapy and pharmacy called to Room 1
    - EM physician assigns roles
    - Airway equipment readied
    - Belmont rapid infuser prepared for IVF 360ml (20ml/kg)
- 00:54 Arrival time
  - HR: 168 RR:58 BP: 78/32 Pox: 87% on NRB
  - Bigamy noted on monitor
  - Pt coughing up blood.
- 00:57 18g IV attempted Right upper ext-Blows
- 00:58 20g IV left hand w/out success
- 00:59 IO placed in right proximal tibia



Children's Hospital  
Colorado

# Case#4 in ED


- 01:00 IVF bolus started via rapid infuser
- 01:02 20g IV in right hand
  - Labs:
    - VBG
    - Blood culture
    - CMP
    - PT/PTT
    - Lactate
- 01:08 IVF bolus 20ml/kg completed
  - HR: 149 BP: 82/32 CR>4 sec RR:64 T:40.4
- 01:10 IVF bolus#2 started on push pull system
- 01:12 Abx ordered/started:
  - Ceftriaxone 100mg/kg
  - Vancomycin 20mg/kg
  - Tamiflu-via NG
- 01:15 Decision made to intubate




Children's Hospital  
Colorado

# Case#4 in ED

- 01:28 2<sup>nd</sup> IVF bolus complete
  - HR: 132 BP: 88/42 Cap Refill~3 sec
- 3<sup>rd</sup> IVF bolus started
- 01:40 Decision made to intubate (HR:138 BP 90/48)
  - RSI and Low Dose Epi (1/10 code dose)
    - Ketamine 2mg/kg
    - Rocuronium 0.5mg/kg
  - Frank blood below cords
- 01:42 Labs:
  - VBG: 7.08/68/10/BE-16
  - CBC: WBC=1.1 ANC=490
  - Lactate: 6.25
  - Glucose: 46
- 01:44 Portable CXR



# 5 y.o. with cough, fever, SOB





# Case#4 5yo fever/cough

- 01:52 5ml/kg D10 given and Norepi drip started
- 01:56 3<sup>rd</sup> IVF bouls completed (total 60 ml/kg)
  - HR 125 BP: 98/46 Cap refill~3 sec
- 02:12 Vitals prior to transfer to ICU
  - HR: 138 BP: 92/48 Cap refill~3 sec




Children's Hospital  
Colorado

# Case#4 5yo in ICU

- 02:30 Repeat Lactate-2.2 (was 6.25)
- 02:40 Left subclavian central line placed
- 07:00 Labs:
  - Influenza B positive
  - Group A Strep positive blood culture

# 6 hours later in PICU

## 5 y.o. with cough, fever, SOB







Children's Hospital  
Colorado

# 5 y.o. with cough, fever, SOB 12 days later

- Pt discharged home and walked out of hospital

**Victory is Mine!**





Children's Hospital  
Colorado

# Important Differences in Cases?

- Similar age patients
  - Similar presentations
  - Same bacteria sepsis
- 
- Why did one patient die and the other walk out of the hospital?



Children's Hospital  
Colorado

# Sepsis Take Home

- The first hour of care saved a life
- EMS recognition and stating concern for sepsis
- Recognition of SHOCK and INFECTION
  - Lactate levels can be helpful
- Access, oxygen, fluid, antibiotics
- Hypotension is bad
- Sepsis protocols work
  - Establishing sepsis clinical care guidelines improves outcomes



# Questions?



**Children's Hospital Colorado**

Affiliated with



University of Colorado  
Anschutz Medical Campus  
School of Medicine

Patrick Mahar, MD

[Patrick.mahar@childrenscolorado.org](mailto:Patrick.mahar@childrenscolorado.org)



**Figure 2. Prevalence of Severity Measures by Categorical Lactate Group**




Figure 2: Prevalence of Severity Measures in Categorical Lactate Groups. Chi-squared test was used to assess for significance of associations.



# SHOCK WARM vs COLD

	WARM shock	COLD shock
<i>Peripheries</i>	warm, flushed	cold, clammy, cyanotic
<i>Capillary refill</i>	< 2 sec	> 2 sec
<i>Pulse</i>	bounding	weak, feeble
<i>Heart rate</i>	tachycardia	tachycardia or bradycardia
<i>Blood pressure</i>	relatively maintained	hypotension
<i>Pulse pressure</i>	widened	narrowed