

Pediatric Assessment

Kelly Reichert, MS RN CCRN
Professional Development Specialist
Outreach Education



Disclosure

Planners, faculty, and others in control of content (either individually or as a group) have no relevant financial relationships with ineligible companies.



Affiliated with
University of Colorado
Anschutz Medical Campus

Objectives



Summarize the differences of pediatric vs adult physiology impacting patient assessment and triage



Develop systematic approach to pediatric assessment and triage



Report escalation of pediatric emergencies and knowledge of resources



Discuss strategies to better support pediatric patients during medical care



Kids are not small adults!



Adults



Kids




Pediatrics vs Adults

Differences	Implications
Greater BSA to body weight	Prone to hypothermia, dehydration
Higher metabolic rate	Prone to hypoxia, hypoglycemia
Higher fluid requirements	Prone to dehydration
Good vasculature and heart	Don't see hypotension until late
Babies are nose breathers	Can't breathe with secretions
Thin chest wall, weak IC muscles	Takes more effort to breathe



Sick or Not Sick: Pediatric Assessment Triangle



Affiliated with
University of Colorado
Anschutz Medical Campus

Appearance - TICLS

TONE - moving? limp?

INTERACTIVENESS - alert? not following?

CONSOLABILITY - by caregiver?

LOOK/GAZE - observant?

SPEECH/CRY - high pitched, hoarse, muffled?





Photo: Children's Hospital Colorado







Affiliated with
University of Colorado
Anschutz Medical Campus

Breathing

Position - extended neck or tripod?

Nasal flaring

Tracheal tug

Head bobbing

Retractions or abdominal breathing


Audible breath sounds: wheeze, stridor, grunting



Photo: Pearson



Photo: Stanford Medicine



Normal



Trouble breathing




Photo: Up To Date



Affiliated with
University of Colorado
Anschutz Medical Campus



Slow RR, Abnormal Positioning
OR Sternal Retractions







Circulation

Skin color:

Early: Pale


Late: Mottled, Cyanotic



Photo: BMJ



Photo: ResearchGate



Mucous membranes

Bruising or purpura?

Mental status – LOC changes



Photo: GrepMed



Photo: Consultant 360



Putting together the ABCs



Concern(s)	Meaning	Interventions
Breathing	Respiratory Distress	Positioning, oxygen, suction, consider meds as applicable
Appearance + Breathing	Respiratory Failure	Positioning, oxygen (mask), suction, get advanced airway ready
Appearance + Circulation	Shock	Oxygen (NRB), access (IV/IO), labs, fluids, reduce oxygen demand / treat cause of shock
Appearance	Neurologic / Metabolic	Check blood sugar, labs, oxygen as needed, determine and treat cause
All 3 (ABC)	Cardiorespiratory Failure	Position, oxygen (bag mask), advanced airway, compressions



Respiratory Distress

Tachypnea
↑ Respiratory Effort
Abnormal Airway Sounds
Retractions
Accessory muscle use
Abdominal breathing

Can quickly progress

Cardiorespiratory Failure

Early

Tachycardia, cool/pale,
decrease UOP

Late

**Bradycardia, hypotension,
cyanosis, unresponsive**

Respiratory Failure



Marked Tachypnea (early)
Apnea (late)
↑/↓ respiratory effort
Poor/absent distal air
Movement
See saw breathing
Tracheal tug
Grunting
Nasal Flaring
Position of comfort

And then...



Affiliated with
University of Colorado
Anschutz Medical Campus



What's Next?

Length-Based Tape

RED to head

Kilogram (kg) weight in pediatrics

- Medication doses
- Equipment



Photo: Armstrong Medical



Weight	Laryngoscope Blade	ET Tube (mm) +0.5 uncuffed	ET Tube Depth (cm)	Suction Cath. (Fr)	IV (ga)	NG (Fr)	Urinary Cath. (Fr)
3-5 kg	1 straight	3.0 Cuffed	3 kg 9-9.5 4 kg 9.5-10 5 kg 10-10.5	8	22-24	5-8	5
6-7 kg	1 Straight	3.0 Cuffed	10.5-11	8	22-24	5-8	8
8-9 kg	1 Straight	3.0 Cuffed	10.5-11	8	22-24	5-8	8
10-11 kg	1 Straight	3.5 Cuffed	11-12	10	20-24	8-10	8-10
12-14 kg	2 Straight	4.0 Cuffed	13.5	10	18-22	10	10
15-18 kg	2 Straight	4.5 Cuffed	14-15	10	18-22	10	10-12
19-23 kg	2 Straight or Curved	5.0 Cuffed	16.5	10	18-20	12-14	10-12
24-29 kg	2 Straight or Curved	6.0 Cuffed	17-18	10	18-20	14-18	12
30-36 kg	3 Straight or Curved	6.5 Cuffed	18.5-19.5	10-12	16-20	16-18	12

PURPLE			
SEIZURE		ICP	
Lorazepam (2 mg/mL) 1 mg (0.5 mL)	3% Saline	21-53 mL	
(4 mg/mL) 1 mg (0.25 mL)	Mannitol (20% 0.2 g/mL)	10 g (50 mL)	
Diazepam IV (5 mg/mL) 2 mg (0.4 mL)	(25% 0.25 g/mL)	10 g (40 mL)	
Phenobarbital (65 mg/mL) 210 mg (3.2 mL)	Furosemide (10 mg/mL)	10 mg (1 mL)	
(130 mg/mL) 210 mg (1.6 mL)	FLUIDS		
Phenytoin (50 mg/mL) 210 mg (4.2 mL)	Fluid Bolus		
Fosphenytoin (50 mg PE/mL) 210 mg PE (4.2 mL)	Crystalloid (NS or LR)	210 mL	
Levetiracetam (100 mg/mL) 625 mg (5.25 mL)	Colloid/Blood	165 mL	
OVERDOSE/HYPOTENSIA		Maintenance	
D ₁₀ W (0.1 g/mL) 5.25 g (52.5 mL)	Dis 1/2 NS + 20 mEq KCl/L		
D ₅ W (0.25 mL) 5.25 g (21 mL)	PAIN		
Naloxone (1 mg/mL) 1 mg (1 mL)	Fentanyl (50 mcg/mL)	10 mcg (0.2 mL)	
(0.4 mg/mL) 1 mg (2.5 mL)	Morphine (2 mg/mL)	1 mg (0.5 mL)	
Flumazenil (0.1 mg/mL) 0.1 mg (1 mL)	(4 mg/mL)	1 mg (0.25 mL)	
Charcoal (25 g/100 mL) 10 g (50 mL)			
Glucagon (1 mg/mL) 0.5 mg (0.5 mL)			
EQUIPMENT		EQUIPMENT	
*E.T. Tube 4.0 Uncuffed/*3.5 Cuffed	Oxygen Mask	Pediatric NRB	
E.T. Insertion Length 11-12 cm	*ETCO ₂	Pediatric	
Styilet 6 French	*Urinary Catheter	8-10 French	
*Suction Catheter 8 French	*Chest Tube	14-20 French	
Laryngoscope 1-1.5 Straight	NG Tube	8-10 French	
BVM Child	Vascular Access	20-24 Ga	
Oral Airway 60 mm	Intraosseous (IO)	15 Ga	
*Nasopharyngeal Airway 18 French	BP Cuff	Child	
*LMA 2	*May not be included in Organizer System(s).		

Photo: HMP Global Learning Network



Affiliated with
University of Colorado
Anschutz Medical Campus

A FULL Set of Vital Signs

Pediatric “normal” varies by age

Think about order!

Heart rate

Respiratory rate - a full minute!

Saturations

Temperature

Blood pressure

CHILDREN'S HOSPITAL COLORADO
OneCall 720-777-3999 | Toll Free 1-800-525-4871

PEDIATRIC VITAL SIGN NORMS

Age	HR Range	Respiratory Rate (Breaths/min)	Blood Pressure		Mean Arterial Pressure (mm Hg)
			Systolic (mm Hg)	Diastolic (mm Hg)	
Birth-28 days	100-205	40-60	67-84	35-53	45-60
1-12 months	100-180	30-53	72-104	37-56	50-62
1-3 yrs.	98-140	22-37	86-106	42-63	49-62
3-5 yrs.	80-120	20-28	89-112	46-72	58-69
5-11 yrs.	75-118	18-25	97-115	57-76	66-72
12-18 yrs.	60-100	12-20	110-131	64-83	73-84

MKTG-6687e@hp-K2.02.14.04



Affiliated with
University of Colorado
Anschutz Medical Campus

Hate the 60s

- Heart Rate **60** = bradycardia
- Respiratory Rate **60** = tachypnea
- Systolic Blood Pressure **60** = hypotension/uncompensated shock
- BGL **60** = borderline
 - 40-60 hypoglycemic



American Heart Association.

AMERICAN ASSOCIATION of CRITICAL CARE NURSES

PALS

Vital Signs in Children

These 3 tables are reproduced or modified from Hazinski MF. Children are different. In: *Nursing Care of the Critically Ill Child*. 3rd ed. Mosby; 2013:1-18, copyright Elsevier.

Normal Heart Rates*

Age	Awake rate	Sleeping rate (beats/min)
Neonate	100-205	90-160
Infant	100-180	90-160
Toddler	98-140	80-120
Preschooler	80-120	65-100
School-age child	75-118	58-90
Adolescent	60-100	50-90

*Always consider the patient's normal range and clinical condition. Heart rate will normally increase with fever or stress.

Normal Respiratory Rates*

Age	Rate (breaths/min)
Infant	30-53
Toddler	22-37
Preschooler	20-28
School-age child	18-25
Adolescent	12-20

*Consider the patient's normal range. The child's respiratory rate is expected to increase in the presence of fever or stress.

Data from Fleming S et al. *Lancet*. 2011;377(9770):1011-1018.



Affiliated with
University of Colorado
Anschutz Medical Campus

Photo: Eastern Illinois University

Pain Scoring

Use the correct tool

- Pain is an abstract concept, hard for kids to number

Kids can withdraw from surroundings when in pain = misunderstandings

Treat pain:

- Fentanyl IV 1mcg/kg
OR Intranasal 2mcg/kg

CRIES -

32-60 weeks gestation

CRIES Scale			
	0	1	2
Crying	None	High-pitched	Inconsolable
Requires O ₂	None	<30% FiO ₂ needed	>30% FiO ₂ needed
Increased vital signs	Normal HR & BP	Increased HR & BP <20%	Increased HR & BP >20%
Expression	Normal	Grimace	Grimace & grunt
Sleeplessness	None	Wakes frequently	Awake constantly

FLACC -

< 3 years or nonverbal

FLACC Score			
CATEGORY	0 POINTS	1 POINT	2 POINTS
Face	Disinterested	Occasional grimace, withdrawn	Frequent frown, clenched jaw
Legs	No position or relaxed	Uneasy, restless, tense	Kicking or legs drawn up
Activity	Normal position	Squirming, tense	Arched, rigid, or jerking
Cry	No crying	Moans or whimpers	Constant crying, screams or sobs
Consolability	Content, relaxed	Distractible	Inconsolable

FACES -

3 - 12 years

Wong-Baker FACES Pain Rating Scale





Systematic Approach



Pediatric Airway Challenges

Same: C-spine stabilization, jaw thrust if needed

Large head, short neck under 2 yo

Larger tongue, larger floppy epiglottis, cricoid cone shaped, small diameter

- “Built-in” obstruction

Infants - obligate nose breathers

- Quick distress with secretions

ABCDE



Photo: Columbia Reports




Photo: Open Anesthesia



Pediatric Airway Interventions

Shoulder Roll / Sniffing Position

- Head tilt chin lift

E-C clamp technique

Suction - use saline!

“Deep suctioning”



Photo: Life with Gremlins

ABCDE



Photo: UCD Emergency Medicine



Photo: Children's Wisconsin



Photo: Anaesthesia, Pain & Intensive Care



Affiliated with
University of Colorado
Anschutz Medical Campus

Pediatric Breathing Challenges

Same: Intubate at GCS 8, significant respiratory failure, compensated shock, laryngeal reflex, impending herniation

Thin chest wall, cartilaginous sternum/ribs, poorly developed intercostal muscles

- Rapid RR, rely on diaphragm/abdominal muscles for respirations

Children have smaller lung capacity and higher oxygen consumption

- Increased RR first sign of distress, hypoxia risk




Photo: Science Direct

Age	Rate
Infant	30-53
Toddler	22-37
Preschooler	20-28
School Aged	18-25
Adolescent	12-20

Source: AHA / PALS



Pediatric Breathing Interventions

Upper vs lower airway

- Stridor vs wheeze

Airway adjuncts

- Bring a bunch to the bedside

Oxygen - cannulas and masks

- NO “blow by”



Photo: Serphinity



Photo: Intersurgical



Photo: Children's Health Ireland



Photo: Flexicare



Photo: Medline



Photo: Grayline

ABCDE

Nasal cannula -

Min: low

Max: Infant 3L,

Pediatric 6L

Simple mask -

Min: 6L

Max: 10L

Non-Rebreather -

Min: 10L (keep bag inflated with breaths)

Max: 15L



Affiliated with
University of Colorado
Anschutz Medical Campus

Pediatric Circulation Challenges

Capillary refill


- More dependent on room-temp

Blood pressure - repeat q15 min

- Kids compensate... hypotension is late

End organ function

- Mental status - parents may notice first!
- Ask about wet diapers



Age	Systolic BP
Term Neonate (0-28 days)	<60
Infants (1-12 months)	<70
Children (1-10 years)	<70 + (age in years x 2)
Children > 10 years	<90

Source: AHA / PALS



Pediatric Circulation Interventions



IVs:
Saphenous
AC/hand
Head



IOs:
Stabilize leg, slight
external rotation
Proximal tibia:
tibial tuberosity
flat, 1-3cm below
knee joint



**Rapid Fluid
Administration:**
3-way connector
Pull/push: pull from
IVF bag into syringe,
push in from syringe



Pediatric Shock

Identification - 3 things at once

- Oxygen - NRB
- Monitor - keep cycling BP!
- Access - IV/IO, don't delay

20 ml/kg isotonic boluses - load them up

- Hepatomegaly, rales/crackles - go slower

“Do not delay inotropes” - PIV, double up

- Epinephrine
- Norepinephrine
- Dopamine

Consider hydrocortisone

Compensated: normotensive

Uncompensated: AMS, hypotension



0 min

Recognize decreased mental status and perfusion.
Begin high flow O₂ and establish IO/IV access according to PALS.

5 min

If no hepatomegaly or rales / crackles then push 20 mL/kg isotonic saline boluses and reassess after each bolus up to 60 mL/kg until improved perfusion. Stop for rales, crackles or hepatomegaly. Correct hypoglycemia and hypocalcemia.
Begin antibiotics.

15 min

Fluid refractory shock?

Begin peripheral IV/IO inotrope infusion, preferably Epinephrine 0.05 – 0.3 µg/kg/min
Use Atropine / Ketamine IV/IO/IM if needed for Central Vein or Airway Access

Titrate Epinephrine 0.05 – 0.3 µg/kg/min for Cold Shock.
(Titrate central Dopamine 5 – 9 µg/kg/min if Epinephrine not available)
Titrate central Norepinephrine from 0.05 µg/kg/min and upward to reverse Warm Shock.
(Titrate Central Dopamine ≥ 10 µg/kg/min if Norepinephrine not available)

60 min

Catecholamine-resistant shock?

If at risk for Absolute Adrenal Insufficiency consider Hydrocortisone.
Use Doppler US, PICCO, FATD or PAC to Direct Fluid, Inotrope, Vasopressor, Vasodilators
Goal is normal MAP-CVP, ScvO₂ > 70%* and CI 3.3 – 6.0 L/min/m²

Algorithm of management of shock in infants and children by American College of Critical Care Medicine

Compensated Shock



Possibly hours

Hypotensive Shock



Potentially minutes

Cardiac Arrest

Pediatric Disability – Da Brain, Dextrose

Same: Mental status - Awake Verbal Pain Unresponsive

Fontanelles! (6-18 months)

Mental status - what's normal?

- You may need the parents

Dextrose - less liver capacity to store glycogen

- Increased risk for hypoglycemia

Head trauma - majority of pediatric trauma deaths

Think about ingestion!

Modified Glasgow Coma Scale for Infants and Children

	Child	Infant	Score
Eye opening	Spontaneous	Spontaneous	4
	To speech	To speech	3
	To pain only	To pain only	2
	No response	No response	1
Best verbal response	Oriented, appropriate	Coos and babbles	5
	Confused	Irritable cries	4
	Inappropriate words	Cries to pain	3
	Incomprehensible sounds	Moans to pain	2
	No response	No response	1
Best motor response*	Obeys commands	Moves spontaneously and purposefully	6
	Localizes painful stimulus	Withdraws to touch	5
	Withdraws in response to pain	Withdraws to response in pain	4
	Flexion in response to pain	Abnormal flexion posture to pain	3
	Extension in response to pain	Abnormal extension posture to pain	2
	No response	No response	1



Pediatric Disability Interventions

Check GLUCOSE!

Rule of 50s:

50 = D50 x 1ml/kg

50 = D25 x 2ml/kg

50 = D10 x 5ml/kg

50 = D5 x 10ml/kg

Suspected Increased ICP:

Head midline, elevate 30

Maintain normotension

Do not excessively hyperventilate - EtCO₂ 35

Consider Mannitol, Hypertonic

Sedation




Photo: Medscape



Photo: CHOP



Pediatric Exposure Interventions

Clothes off - look under diaper too

- Then bundle them back up

Aim for normothermia

Warm: Blankets, Warmed IVF, Bair Hugger

Cool: Remove layers, Wet washcloths,
Cooling blankets, antipyretics



Photo: Wyoming Department of Health





Supporting Pediatric Patients

“We owe it to the future not to harm our children in their hearts and minds while we cure their diseases and repair their broken bones.”

Pate, JT et al. (1996)

Comfort Holds

Snuggle and Swaddle



YouTube

Back to Chest



Chest to Chest



<https://www.youtube.com/watch?v=6b-CV6kXgVA>



Affiliated with
University of Colorado
Anschutz Medical Campus

How to Use Comfort Holds During a Procedure

Children's Hospital Colorado
123K subscribers

Subscribe

46 Share Download Clip Save

Child Development Considerations

- Younger children
 - Separation anxiety
 - Involve caregivers (and patient as able)
 - Toddlers - offer choices
- School age children
 - Fear loss of competence or control
 - Involve the patient - helpers, writers, give them a job!
- Teenagers
 - Vague in complaints and needs
 - Fear being different
 - Normalize experience



Preparation using their senses

**Some kids say it sounds like...
feels like... smells like...**

See:

- Soft straw (IV)
- Bright lights (exam lights)

Hear:

- Loud noises like construction (MRI)
- Popping like a soda can opening (J-tip)

Taste:

- Salty (saline)
- Sprite without bubbles (oral contrast)

Feel:

- Cold, wet (soap)
- Tight hug/squeeze (tourniquet)
- Quick pinch/poke (IV)

Smell:

- The ocean (saline)
- Hand sanitizer (Chloraprep)



Child Development Considerations

AVOID:

“Don’t move while I do this”

“The IV will hurt”

“It will burn”

“It will taste bad”

“Show me how brave you are / what a big kid you are”

TRY:

“Your job is to hold as still as you can”

“You’ll feel a pinch/poke”

“It might feel warm / cool going in”

“It might taste bitter”

“Remember, your job is to be as still as you can. It’s OK to cry. I know this is scary.”



Alternative Focus / Distraction

- Tablet / smartphone
- Search and Find Books
- Stress balls
- Pinwheels
- Music / singing
- Deep breathing
- Grounding activities - counting



PANDA UP

P = Prepare *Use prep supplies and treatment room, educate family*

A = Anxiety Reduction *Implement relaxation methods and coping plan*

N = Numb *Use numbing agents prior to procedures; sucrose for infants*

D = Distract *Apply methods such as vibration tool and alternative focus*

A = Attitude *Maintain a calm, positive attitude*

U = Use One Person's Voice *Understand everyone's role*

P = Position *Use comfort positioning*



Take Away Points

- Keep in mind pediatric differences
- Use validated tools and a systematic approach
- Basic interventions save lives
- Early recognition of decompensation is key
- PANDA UP for procedures
- Know your resources:

CHCO OneCall 720-777-3999

CHCO Pathways

<https://www.childrenscolorado.org/health-professionals/clinical-resources/clinical-pathways/>



Affiliated with
University of Colorado
Anschutz Medical Campus

Thank you Questions?

