

AGES 0–4 YEARS: STEPWISE APPROACH FOR MANAGEMENT OF ASTHMA

	Intermittent Asthma	Management of Persistent Asthma in Individuals Ages 0–4 Years				
Treatment	STEP 1	STEP 2	STEP 3	STEP 4	STEP 5	STEP 6
Preferred	PRN SABA and At the start of RTI: Add short course daily ICS▲	Daily low-dose ICS and PRN SABA	Daily low-dose ICS-LABA and PRN SABA▲ or Daily low-dose ICS + montelukast,* or daily medium-dose ICS, and PRN SABA	Daily medium-dose ICS-LABA and PRN SABA	Daily high-dose ICS-LABA and PRN SABA	Daily high-dose ICS-LABA + oral systemic corticosteroid and PRN SABA
Alternative		Daily montelukast* or Cromolyn,* and PRN SABA		Daily medium-dose ICS + montelukast* and PRN SABA	Daily high-dose ICS + montelukast* and PRN SABA	Daily high-dose ICS + montelukast*+ oral systemic corticosteroid and PRN SABA
For children age 4 years only, see Step 3 and Step 4 on Management of Persistent Asthma in Individuals Ages 5–11 Years diagram.						
Assess Control <ul style="list-style-type: none"> First check adherence, inhaler technique, environmental factors, ▲ and comorbid conditions. Step up if needed; reassess in 4–6 weeks Step down if possible (if asthma is well controlled for at least 3 consecutive months) Consult with asthma specialist if Step 3 or higher is required. Consider consultation at Step 2. <p>Control assessment is a key element of asthma care. This involves both impairment and risk. Use of objective measures, self-reported control, and health care utilization are complementary and should be employed on an ongoing basis, depending on the individual's clinical situation.</p>						

Abbreviations: ICS, inhaled corticosteroid; LABA, long-acting beta₂-agonist; SABA, inhaled short-acting beta₂-agonist; RTI, respiratory tract infection; PRN, as needed

▲ Updated based on the 2020 guidelines.

* Cromolyn and montelukast were not considered for this update and/or have limited availability for use in the United States. The FDA issued a Boxed Warning for montelukast in March 2020.

NOTES FOR INDIVIDUALS AGES 0–4 YEARS DIAGRAM

Quick-relief medications

- Use SABA as needed for symptoms. The intensity of treatment depends on severity of symptoms: up to 3 treatments at 20-minute intervals as needed.
- **Caution:** Increasing use of SABA or use >2 days a week for symptom relief (not prevention of EIB) generally indicates inadequate control and may require a step up in treatment.
- Consider short course of oral systemic corticosteroid if exacerbation is severe or individual has history of previous severe exacerbations.

Each step: Assess environmental factors, provide patient education, and manage comorbidities▲

- In individuals with sensitization (or symptoms) related to exposure to pests‡: conditionally recommend integrated pest management as a single or multicomponent allergen-specific mitigation intervention.▲
- In individuals with sensitization (or symptoms) related to exposure to identified indoor allergens, conditionally recommend a multi-component allergen-specific mitigation strategy.▲
- In individuals with sensitization (or symptoms) related to exposure to dust mites, conditionally recommend impermeable pillow/mattress covers only as part of a multicomponent allergen-specific mitigation intervention, but not as a single component intervention.▲

Notes

- If clear benefit is not observed within 4–6 weeks and the medication technique and adherence are satisfactory, the clinician should consider adjusting therapy or alternative diagnoses.

Abbreviations

EIB, exercise-induced bronchoconstriction; SABA, inhaled short-acting beta₂-agonist.
 ▲Updated based on the 2020 guidelines.
 ‡ Refers to mice and cockroaches, which were specifically examined in the Agency for Healthcare Research and Quality systematic review.

AGES 5-11 YEARS: STEPWISE APPROACH FOR MANAGEMENT OF ASTHMA

	Intermittent Asthma	Management of Persistent Asthma in Individuals Ages 5-11 Years				
Treatment	STEP 1	STEP 2	STEP 3	STEP 4	STEP 5	STEP 6
Preferred	PRN SABA	Daily low-dose ICS and PRN SABA	Daily and PRN combination low-dose ICS-formoterol▲	Daily and PRN combination medium-dose ICS-formoterol▲	Daily high-dose ICS-LABA and PRN SABA	Daily high-dose ICS-LABA + oral systemic corticosteroid and PRN SABA
Alternative		Daily LTRA,* or Cromolyn,* or Nedocromil,* or Theophylline,* and PRN SABA	Daily medium-dose ICS and PRN SABA or Daily low-dose ICS-LABA, or daily low-dose ICS + LTRA,* or daily low-dose ICS +Theophylline,* and PRN SABA	Daily medium-dose ICS-LABA and PRN SABA or Daily medium-dose ICS + LTRA* or daily medium-dose ICS + Theophylline,* and PRN SABA	Daily high-dose ICS + LTRA* or daily high-dose ICS + Theophylline,* and PRN SABA	Daily high-dose ICS + LTRA* + oral systemic corticosteroid or daily high-dose ICS + Theophylline* + oral systemic corticosteroid, and PRN SABA
		Steps 2-4: Conditionally recommend the use of subcutaneous immunotherapy as an adjunct treatment to standard pharmacotherapy in individuals > 5 years of age whose asthma is controlled at the initiation, build up, and maintenance phases of immunotherapy▲			Consider Omalizumab**▲	

Assess Control

- First check adherence, inhaler technique, environmental factors,▲ and comorbid conditions.
- Step up** if needed; reassess in 2-6 weeks
- Step down** if possible (if asthma is well controlled for at least 3 consecutive months)

Consult with asthma specialist if Step 4 or higher is required. Consider consultation at Step 3.

Control assessment is a key element of asthma care. This involves both impairment and risk. Use of objective measures, self-reported control, and health care utilization are complementary and should be employed on an ongoing basis, depending on the individual's clinical situation.

Abbreviations: ICS, inhaled corticosteroid; LABA, long-acting beta₂-agonist; LTRA, leukotriene receptor antagonist; SABA, inhaled short-acting beta₂-agonist

[▲] Updated based on the 2020 guidelines.

* Cromolyn, Nedocromil, LTRAs including montelukast, and Theophylline were not considered in this update and/or have limited availability for use in the United States, and/or have an increased risk of adverse consequences and need for monitoring that make their use less desirable. The FDA issued a Boxed Warning for montelukast in March 2020.

** Omalizumab is the only asthma biologic currently FDA-approved for this age range.



NOTES FOR INDIVIDUALS AGES 5–11 YEARS DIAGRAM

Quick-relief medications	<ul style="list-style-type: none"> • Use SABA as needed for symptoms. The intensity of treatment depends on severity of symptoms: up to 3 treatments at 20-minute intervals as needed. • In Steps 3 and 4, the preferred option includes the use of ICS-formoterol 1 to 2 puffs as needed up to a maximum total daily maintenance and rescue dose of 8 puffs (36 mcg).▲ • Caution: Increasing use of SABA or use >2 days a week for symptom relief (not prevention of EIB) generally indicates inadequate control and may require a step up in treatment.
Each step: Assess environmental factors, provide patient education, and manage	<ul style="list-style-type: none"> • In individuals with sensitization (or symptoms) related to exposure to pests‡: conditionally recommend integrated pest management as a single or multicomponent allergen-specific mitigation intervention.▲ • In individuals with sensitization (or symptoms) related to exposure to identified indoor allergens, conditionally recommend a multi-component allergen-specific mitigation strategy.▲ • In individuals with sensitization (or symptoms) related to exposure to dust mites, conditionally recommend impermeable pillow/mattress covers only as part of a multicomponent allergen-specific mitigation intervention, but not as a single component intervention.▲
Notes	<ul style="list-style-type: none"> • The terms ICS-LABA and ICS-formoterol indicate combination therapy with both an ICS and a LABA, usually and preferably in a single inhaler. • Where formoterol is specified in the steps, it is because the evidence is based on studies specific to formoterol. • In individuals ages 5–11 years with persistent allergic asthma in which there is uncertainty in choosing, monitoring, or adjusting anti-inflammatory therapies based on history, clinical findings, and spirometry, FeNO measurement is conditionally recommended as part of an ongoing asthma monitoring and management strategy that includes frequent assessment.
Abbreviations	<p>2</p> <p>EIB (exercise-induced bronchoconstriction); FeNO (fractional exhaled nitric oxide); ICS (inhaled corticosteroid); LABA (long-acting beta -agonist); SABA (inhaled short-acting beta -agonist). ▲Updated based on the 2020 guidelines. ‡ Refers to mice and cockroaches, which were specifically examined in the Agency for Healthcare Research and Quality systematic review.</p>

AGES 12+ YEARS: STEPWISE APPROACH FOR MANAGEMENT OF ASTHMA

	Intermittent Asthma	Management of Persistent Asthma in Individuals Ages 12+ Years				
Treatment	STEP 1	STEP 2	STEP 3	STEP 4	STEP 5	STEP 6 [■]
Preferred	PRN SABA	Daily low-dose ICS and PRN SABA or PRN concomitant ICS and SABA▲	Daily and PRN combination low-dose ICS-formoterol▲	Daily and PRN combination medium-dose ICS-formoterol▲	Daily medium-high dose ICS-LABA + LAMA and PRN SABA▲	Daily high-dose ICS-LABA + oral systemic corticosteroids + PRN SABA
Alternative		Daily LTRA* and PRN SABA or Cromolyn,* or Nedocromil,* or Zileuton,* or Theophylline,* and PRN SABA	Daily medium-dose ICS and PRN SABA or Daily low-dose ICS-LABA, or daily low-dose ICS + LAMA,▲ or daily low-dose ICS + LTRA,* and PRN SABA or Daily low-dose ICS + Theophylline* or Zileuton,* and PRN SABA	Daily medium-dose ICS-LABA or daily medium-dose ICS + LAMA, and PRN SABA▲ or Daily medium-dose ICS + LTRA,* or daily medium-dose ICS + Theophylline,* or daily medium-dose ICS + Zileuton,* and PRN SABA	Daily medium-high dose ICS-LABA or daily high-dose ICS + LTRA,* and PRN SABA	
		Steps 2–4: Conditionally recommend the use of subcutaneous immunotherapy as an adjunct treatment to standard pharmacotherapy in individuals ≥ 5 years of age whose asthma is controlled at the initiation, build up, and maintenance phases of immunotherapy▲			Consider adding Asthma Biologics (e.g., anti-IgE, anti-IL5, anti-IL5R, anti-IL4/IL13)**	

Assess Control

- First check adherence, inhaler technique, environmental factors,▲ and comorbid conditions.
- Step up** if needed; reassess in 2–6 weeks
- Step down** if possible (if asthma is well controlled for at least 3 consecutive months)

Consult with asthma specialist if Step 4 or higher is required. Consider consultation at Step 3.

Control assessment is a key element of asthma care. This involves both impairment and risk. Use of objective measures, self-reported control, and health care utilization are complementary and should be employed on an ongoing basis, depending on the individual's clinical situation.

Abbreviations: ICS, inhaled corticosteroid; LABA, long-acting beta₂-agonist; LAMA, long-acting muscarinic antagonist; LTRA, leukotriene receptor antagonist; SABA, inhaled short-acting beta₂-agonist

[▲] Updated based on the 2020 guidelines.

* Cromolyn, Nedocromil, LTRAs including Zileuton and montelukast, and Theophylline were not considered for this update, and/or have limited availability for use in the United States, and/or have an increased risk of adverse consequences and need for monitoring that make their use less desirable. The FDA issued a Boxed Warning for montelukast in March 2020.

** The AHRQ systematic reviews that informed this report did not include studies that examined the role of asthma biologics (e.g. anti-IgE, anti-IL5, anti-IL5R, anti-IL4/IL13). Thus, this report does not contain specific recommendations for the use of biologics in asthma in Steps 5 and 6.

■ Data on the use of LAMA therapy in individuals with severe persistent asthma (Step 6) were not included in the AHRQ systematic review and thus no recommendation is made.



NOTES FOR INDIVIDUALS AGES 12+ YEARS DIAGRAM

Quick-relief medications	<ul style="list-style-type: none"> • Use SABA as needed for symptoms. The intensity of treatment depends on the severity of symptoms: up to 3 treatments at 20-minute intervals as needed. • In steps 3 and 4, the preferred option includes the use of ICS-formoterol 1 to 2 puffs as needed up to a maximum total daily maintenance and rescue dose of 12 puffs (54 mcg).▲ • Caution: Increasing use of SABA or use >2 days a week for symptom relief (not prevention of EIB) generally indicates inadequate control and may require a step up in treatment.
Each step: Assess environmental factors, provide patient education, and manage	<ul style="list-style-type: none"> • In individuals with sensitization (or symptoms) related to exposure to pests‡: conditionally recommend integrated pest management as a single or multicomponent allergen-specific mitigation intervention.▲ • In individuals with sensitization (or symptoms) related to exposure to identified indoor allergens, conditionally recommend a multi-component allergen-specific mitigation strategy.▲ • In individuals with sensitization (or symptoms) related to exposure to dust mites, conditionally recommend impermeable pillow/mattress covers only as part of a multicomponent allergen-specific mitigation intervention, but not as a single component intervention.▲
Notes	<ul style="list-style-type: none"> • The terms ICS-LABA and ICS-formoterol indicate combination therapy with both an ICS and a LABA, usually and preferably in a single inhaler. • Where formoterol is specified in the steps, it is because the evidence is based on studies specific to formoterol. • In individuals ages 12 years and older with persistent allergic asthma in which there is uncertainty in choosing, monitoring, or adjusting anti-inflammatory therapies based on history, clinical findings, and spirometry, FeNO measurement is conditionally recommended as part of an ongoing asthma monitoring and management strategy that includes frequent assessment. • Bronchial thermoplasty was evaluated in Step 6. The outcome was a conditional recommendation against the therapy.
Abbreviations 2	<p>EIB, exercise-induced bronchoconstriction; FeNO, fractional exhaled nitric oxide; ICS, inhaled corticosteroid; LABA, long-acting beta -agonist; SABA, inhaled short-acting beta -agonist.</p> <p>▲Updated based on the 2020 guidelines.</p> <p>‡ Refers to mice and cockroaches, which were specifically examined in the Agency for Healthcare Research and Quality systematic review.</p>